

425x160mm



Gel Cold Pack
User Manual

MODEL: REX162pack1-XL

Important:
Read carefully and keep these instructions for future reference.

Caution

Do not use on broken skin, open wounds, severe swelling, or sensitive skin.

Avoid prolonged use. Do not use for long periods as it can cause frostbite.

Do not heat the pack in the microwave as it can damage the product.

Keep out of reach of children.
If the pack tears or breaks, discard immediately. **DO NOT use** on people unable to remove the product, including children, infants, and elderly, or on people unable to follow all use instructions.

Stop use if there are any blisters, red marks, or adverse reactions to the cold pack. **Ask a doctor before use** if you have diabetes, poor circulation or heart disease, rheumatoid arthritis, or are pregnant.

Instructions for Use

Preparation:

Place the gel pack in the included storage bag in the freezer for at least 4 hours before use.

Keeping the ice pack stored in the freezer will ensure it's always ready for emergency use.

Application:

- 1. Lay the pack onto the injury area with the soft side facing the skin.
- 2. Wrap the straps around the limb, pulling to the preferred amount of compression.
- 3. Use for up to 20 minutes every 2 to 4 hours as needed.



Freezing Time (0°F / -18°C)	Lasts for about
Less than 4 hours	10 minutes*
4-6 hours	20 minutes*
Over 6 hours	30 minutes*

*Lasting times may vary due to surrounding temperatures.

Recommendation of Use.

Use the plush side for 10 min, then use the nylon inner for another 10 min. If there is condensation water on the nylon side, just wipe it off.

Warranty

The REVIX gel cold pack is warranted against defects in materials and workmanship for a lifetime from the date of purchase. The warranty does not cover damage from misuse, accidents, alterations, or disassembly.

Need Help?

For any questions, contact REVIX customer support at support@revixcare.com.

